



Document allergies on approved form and ensure medication reconciliation has been reviewed as per organizational process

Safe initiation of DOAC for patients with CAD/PAD already on guideline directed therapy

ACTION

Administration

DOCUMENT PURPOSE

This order set may be used for patients diagnosed with stable coronary artery disease (CAD) and/or peripheral artery disease (PAD) who are already being treated with guideline directed antiplatelet therapy and require prescription of adjunct direct oral anticoagulant therapy (currently only rivaroxaban) in an inpatient or outpatient setting.

Rivaroxaban is indicated, in combination with aspirin, to reduce the risk of major cardiovascular events (cardiovascular [CV] death, myocardial infarction [MI], and stroke) in patients with chronic CAD or PAD ⁽¹⁾.

Other FDA approved pharmacological therapies for cardiovascular prevention can include ^(1,2,3):

- Beta blockers
- Blood pressure lowering medication
- Glucose lowering medication
- LDL cholesterol lowering medication
 - Statins
 - PCSK9 inhibitors
 - Other lipid lowering therapies
- Renin angiotensin blockers (ACE-I/ARB)
- Antithrombotic therapies

Note: Specific use of these medications are not addressed in this order set

Eligibility

INCLUSION CRITERIA

Patients should have the following to be eligible for rivaroxaban + aspirin as a secondary prevention strategy ⁽¹⁾:

CORONARY ARTERY DISEASE (CAD) INDICATIONS (SELECT AT LEAST 1 INDICATION AND 1 HIGH-RISK FEATURE)

(1) DOCUMENTED CAD

- Prior myocardial infarction, or
- Documented multi-vessel coronary disease (by angiography, stress testing, or coronary revascularization) with a history of stable or unstable angina, or
- Multi-vessel percutaneous coronary intervention (PCI), or
- Multi-vessel coronary artery bypass graft (CABG) surgery

Submitted by:	_____	_____	_____	<input type="checkbox"/> Read Back
	ID	PRINTED NAME	YYYY-MM-DD HH:MM	
Practitioner:	_____	_____	_____	_____
	ID	PRINTED NAME	YYYY-MM-DD HH:MM	SIGNATURE



Document allergies on approved form and ensure medication reconciliation has been reviewed as per organizational process

Safe initiation of DOAC for patients with CAD/PAD already on guideline directed therapy ACTION

Eligibility Continued

INCLUSION CRITERIA CONTINUED

(2) HIGH-RISK FEATURES

Subjects with CAD must also meet at least one of the following criteria:

- Age ≥65, or
- Age <65 and other high-risk features
 - Documented atherosclerosis or revascularization involving at least 2 vascular beds, OR
 - At least 2 additional risk factors:
 - Current or recent smoker (quit within 1 year)
 - Diabetes mellitus
 - Renal dysfunction with estimated glomerular filtration rate <60 ml/min
 - History of symptomatic heart failure
 - Non-lacunar ischemic stroke ≥1 month ago

AND/OR

PERIPHERAL ARTERY DISEASE (PAD), SELECT AT LEAST 1

- Previous PAD revascularization (e.g. aorto-femoral bypass surgery, limb bypass surgery, or percutaneous transluminal angioplasty revascularization of the iliac, or infra-inguinal arteries), or
- Previous limb or foot amputation for arterial vascular disease, or
- History of intermittent claudication and one or more of the following:
 - An ankle-brachial index (ABI) < 0.90, or
 - Significant peripheral artery stenosis (≥50%) documented by angiography, or by duplex ultrasound, or
- Significant carotid artery disease (previous carotid revascularization or asymptomatic carotid artery stenosis ≥50%).

Note: Patients with severe heart failure (ejection fraction <30% or New York Heart Association class III or IV) were not included in the COMPASS randomized trial.

Reference Document Only
© 2018 ACForum. All rights reserved. Unauthorized use, reproduction or disclosure is prohibited.

Submitted by: _____ Read Back

ID PRINTED NAME YYYY-MM-DD HH:MM

Practitioner: _____

ID PRINTED NAME YYYY-MM-DD HH:MM SIGNATURE



Document allergies on approved form and ensure medication reconciliation has been reviewed as per organizational process

Safe initiation of DOAC for patients with CAD/PAD already on guideline directed therapy

ACTION

Eligibility Continued

USE NOT RECOMMENDED

The following criteria indicate scenarios where rivaroxaban + aspirin should NOT be used as a secondary cardiovascular prevention strategy⁽¹⁾:

- Active pathological bleeding
- Advanced chronic kidney disease (estimated CrCl <15 ml per minute)
- Mechanical heart valves
- Other indication for anticoagulation (e.g. atrial fibrillation or venous thromboembolism) or antiplatelet medication (e.g. cilostazole)
- Current need for dual antiplatelet therapy (DAPT): aspirin plus P2Y12 antiplatelet medicine (clopidogrel, prasugrel, or ticagrelor)
- Pregnancy/breastfeeding
- Severe hypersensitivity reaction to rivaroxaban or aspirin (e.g. anaphylactic reactions)
- Any stroke within 1 month
- Systemic treatment with strong inhibitors of both CYP 3A4 and p-glycoprotein or strong inducers of CYP 3A4

USE WITH CAUTION

The following criteria indicate scenarios where use of rivaroxaban + aspirin should be done after carefully considering risks and benefits:

- Any history of pathologic bleeding
- High risk of pathologic bleeding
- History of hemorrhagic or lacunar stroke (>1 month)

Reassess candidacy for anticoagulant prophylaxis as clinically indicated or with any medication changes.

Reference Document Only
© 2018 ACForum. All rights reserved. Unauthorized use, reproduction or disclosure is prohibited.

Submitted by:

Read Back

Practitioner:



Document allergies on approved form and ensure medication reconciliation has been reviewed as per organizational process

Safe initiation of DOAC for patients with CAD/PAD already on guideline directed therapy ACTION

Factors Influencing Drug Selection

Renal and liver characteristics are necessary to determine appropriateness of anticoagulation therapy.

RENAL FUNCTION

Calculate estimated CrCl using the Cockcroft-Gault formula based on the following:

$$\frac{[(140 - \text{Age}) \times \text{actual weight in kg}]}{[72 \times \text{serum creatinine}]} \times 0.85 \text{ if female}$$

Age: _____

Actual body weight: _____ (kg)

Gender: _____

Serum creatinine: _____ (mg/dL)

Estimated CrCl: _____ (mL/minute)

RENAL IMPAIRMENT

AVOID use in patients with an estimated CrCl <15 ml/min

LIVER FUNCTION

Liver disease: No Yes: Child Pugh Grade: _____

CHILD PUGH SCORE

Measure	1 point	2 points	3 points
Total bilirubin (mg/dL)	< 2	2 - 3	> 3
Serum albumin (g/dL)	> 3.5	2.8 - 3.5	< 2.8
INR	Less than 1.7	1.7 – 2.2	Greater than 2.2
Ascites	None	Mild (or suppressed with medication)	Moderate to Severe (or refractory)
Hepatic encephalopathy	None	Grade I-II	Grade III-IV

Note: The score employs five clinical measures of liver disease⁽⁴⁾.

Each measure is scored 1-3, with 3 indicating the worst condition.

Total score of 5-6: grade A (well-compensated disease)

Total score of 7-9: grade B (significant functional compromise)

Total score 10-15: grade C (decompensated disease)

HEPATIC IMPAIRMENT

Child-Pugh A: No adjustment needed

Child-Pugh B/C: **AVOID** use per package insert

Reference Document Only
© 2018 ACForum. All rights reserved. Unauthorized use, reproduction or disclosure is prohibited.

Submitted by: _____ Read Back

ID _____ PRINTED NAME _____ YYYY-MM-DD HH:MM _____

Practitioner: _____

ID _____ PRINTED NAME _____ YYYY-MM-DD HH:MM _____ SIGNATURE _____



Document allergies on approved form and ensure medication reconciliation has been reviewed as per organizational process

Safe initiation of DOAC for patients with CAD/PAD already on guideline directed therapy ACTION

Other Considerations

PROTON PUMP INHIBITORS (PPIs)

Note: Clinician may consider PPI or other form of gastroprotection for patients at high risk of GI bleeding, particularly if using multiple antithrombotic agents or with a prior history of upper GI bleeding ^(6, 7, 8).

MANAGED CARE

Complete prior authorization paperwork if required by payer (<https://www.covermymeds.com>)

SHARED DECISION-MAKING DISCUSSION

Note: If drug costs are a barrier to filling prescriptions for medication, refer patient to appropriate resources.

Select all that have been discussed with patient

- Bleeding risk/reversal agents
- Dosing regimen options (e.g. once vs. twice daily)
- Lifestyle factors of drug (e.g. diet, blood draws, activities)
- Out-of-pocket medication cost discussed with patient
- Other (specify): _____

Patient Education

Provide applicable education materials/instructions to the patient as per policy/procedure ^(1, 6).

The following topics are important to include within patient education:

- Follow-up appointments for blood work
 - Follow-up contact information: _____
- Safety net phone number to call if any barriers or issues after discharge: _____
- Medication management, including starting/stopping new medication, missed doses and dose change (dose de-escalation or switch to oral therapy at appropriate date/time)
- Importance of medication adherence
- Expected duration of anticoagulation therapy
- Drug/diet considerations (if any)
- Bleeding and bruising risks
- When to seek medical attention (e.g. warning signs for bleeding, symptoms of VTE)
- Written education materials for patient/family/caregivers to review after discharge
- Medication reconciliation completed
- Patient education documented per health system policy

Reference Document Only
© 2018 ACForum. All rights reserved. Unauthorized use, reproduction or disclosure is prohibited.

Submitted by: _____ Read Back

ID PRINTED NAME YYYY-MM-DD HH:MM

Practitioner: _____

ID PRINTED NAME YYYY-MM-DD HH:MM SIGNATURE



Document allergies on approved form and ensure medication reconciliation has been reviewed as per organizational process

Safe initiation of DOAC for patients with CAD/PAD already on guideline directed therapy

ACTION

References

1. Eikelboom JW, Connolly SJ, Bosch J, et al. Rivaroxaban with or without Aspirin in Stable Cardiovascular Disease. *New England Journal of Medicine*. 2017;377(14):1319-1330. doi:[10.1056/NEJMoa1709118](https://doi.org/10.1056/NEJMoa1709118).
2. Arnett DK, Goodman RA, Halperin JL, Anderson JL, Parekh AK, Zoghbi WA. AHA/ACC/HHS Strategies to Enhance Application of Clinical Practice Guidelines in Patients with Cardiovascular Disease and Comorbid Conditions. *Journal of the American College of Cardiology*. 2014;64(17):1851-1856. doi:[10.1016/j.jacc.2014.07.012](https://doi.org/10.1016/j.jacc.2014.07.012)
3. Anderson JL, Halperin JL, Albert N, et al. Management of Patients With Peripheral Artery Disease (Compilation of 2005 and 2011 ACCF/AHA Guideline Recommendations): A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*. 2013;61(14):1555-1570. doi:[10.1016/j.jacc.2013.01.004](https://doi.org/10.1016/j.jacc.2013.01.004)
4. Angermayr, B., Cejna, M., Karnel, F., Gschwantler, M., Koenig, F., ... Peck-Radosavljevic, M. (2003). Child Pugh versus MELD score in predicting survival in patients undergoing transjugular intrahepatic portosystemic shunt. *Gut*, 52(6), 879–85. doi:10.1136/gut.52.6.879
5. Michigan Anticoagulation Quality Improvement Initiative (MAQI2). (2017). Anticoagulation toolkit: A consortium-developed quick reference for anticoagulation.
6. Slott, Amy, Tschurtz, Brette, Williams, Scott. Discharge Instructions / Education Materials for Venous Thromboembolism (VTE): A Comprehensive Approach to Medication Management. https://www.jointcommission.org/assets/1/6/FINAL_WEB_VTE_COMPENDIUM_OF_RESOURCE_S_09152017.pdf. Accessed December 19, 2017.
7. Bhagirath VC, Eikelboom JW, Anand SS. Low-dose rivaroxaban plus aspirin for the prevention of cardiovascular events: an evaluation of COMPASS. *Future Cardiol*. 2018;14(6):443-453. doi:[10.2217/fca-2018-0059](https://doi.org/10.2217/fca-2018-0059)
8. Bosch J, Eikelboom JW, Connolly SJ, et al. Rationale, Design and Baseline Characteristics of Participants in the Cardiovascular Outcomes for People Using Anticoagulation Strategies (COMPASS) Trial. *Can J Cardiol*. 2017;33(8):1027-1035. doi:[10.1016/j.cjca.2017.06.001](https://doi.org/10.1016/j.cjca.2017.06.001)

Reference Document Only
© 2018 ACForum. All rights reserved. Unauthorized use, reproduction or disclosure is prohibited.

Submitted by:

ID

PRINTED NAME

YYYY-MM-DD HH:MM

Read Back

Practitioner:

ID

PRINTED NAME

YYYY-MM-DD HH:MM

SIGNATURE